

Neurotic disorders

James Lindesay and Tony Winston

University of Leicester, Leicester, UK

Introduction

This paper updates a previous review of the topic published in this series in 1992.¹ There are no major advances in knowledge, understanding or treatment to report, but there has been a steady growth in interest, articles, chapters and textbooks dealing with neurotic disorders in the elderly, which is an encouraging sign that these common, disabling and treatable conditions are now being taken more seriously by health professionals.

The concept of neurosis

Debate and discussion continue about the relative validity of categorical and dimensional concepts of neurosis. The unitary, dimensional concept of neurosis invented and developed in the eighteenth and nineteenth centuries has yielded in the second half of the twentieth century to a system of specific diagnostic categories based upon patterns of symptoms. This system is currently enshrined in the tenth edition of the *International classification of diseases* (ICD-10)² and the fourth edition of the *Diagnostic and statistical manual of mental disorders* (DSM-IV).³ ICD-10 still makes reference to a group of 'neurotic, stress-related and somatoform disorders', but in DSM-IV all mention of neurosis has been removed. Despite this powerful movement away from the dimensional concept of neurosis in the modern nosologies, the categorical alternative they provide is unsatisfactory in several respects. There is considerable comorbidity between the specific neurotic diagnoses, and between them and other disorders such as depression. These categories are also unstable over time, and the effectiveness of treatments appears to be

relatively independent of diagnosis.^{4,5} The dimensionality of neurotic disorders is particularly apparent in community and primary care populations, where independent but related dimensions of depression and anxiety underlie the manifest psychological symptomatology in both younger and older adults.^{6,7} An individual's symptoms during any particular episode are determined by pre-existing vulnerability, the particular destabilization factors precipitating that episode, and the steps taken by the patient or the doctor to manage it (restitution factors). Most of the current diagnostic categories such as phobic disorder, dissociative disorder or somatoform disorder may be seen as the results of various maladaptive attempts by patients to reduce their symptoms.⁸ The dimensional approach to neurotic disorders formulates them as longitudinal processes rather than just as cross-sectional episodes; this perspective is particularly important in elderly patients who often present with many years of illness experience.⁹ It also integrates the depressive component of many neurotic disorders, something that categorical classifications have never done adequately.

Epidemiology

One important consequence of the modern classificatory approach to neurotic disorders is that specific diagnostic criteria and standardized methods of assessment have made it possible to collect reliable and comparable information about the distribution of neurotic symptoms and disorders in populations, and the various factors affecting this distribution. However, while modern epidemiological methods allow for reliability and comparability, the validity of the various case definitions used remains questionable. Different rules with regard to symptom definition, severity and diagnostic hierarchies result in very different definitions of 'caseness', particularly with neurotic disorders where there is a very wide spectrum of severity.¹⁰

Address for correspondence: Professor James Lindesay, Division of Psychiatry for the Elderly, University of Leicester, Leicester General Hospital, Leicester LE5 4PW, UK.

To a great extent, the prevalence rates of neurotic disorders found in the elderly depend upon the population investigated. They are uncommon primary diagnoses in hospital inpatient, outpatient and casualty populations,¹¹⁻¹³ and while there is a steady accumulation of chronic cases in primary care settings, there is a decline in the rate of new consultations with age.^{14,15} As with younger adults, the highest rates of neurotic disorders in the elderly, particularly anxiety, are found in the community population. To some extent this is because a proportion of the community cases identified by epidemiological surveys are relatively mild, transient or unproblematic, but it is also true that there are clinically significant cases that either do not present to health services or are not identified or treated as such if they do.¹⁶⁻¹⁹

Table 1 summarizes the findings of the United States Epidemiologic Catchment Area (ECA) study so far as DSM-III categories related to neurotic disorder in the elderly are concerned.²⁰⁻²² Overall, lifetime prevalence rates decreased with age, although this decline was least pronounced in phobic disorder and somatization disorder; phobic disorder was the commonest psychiatric

disorder identified in women over 65 years, and the second commonest after cognitive impairment in men. Women had higher period prevalence rates than men for phobic disorder, panic disorder, generalized anxiety disorder and somatization disorder, and higher lifetime prevalence rates of obsessive-compulsive disorder and somatization disorder. Most elderly subjects with neurotic disorders had developed them before their fifties, but elderly cases of phobic disorder, panic disorder and obsessive-compulsive disorder tended to be of later onset. Incidence rates of most neurotic disorders fell with age in both sexes, but this was least apparent for phobic disorder and obsessive-compulsive disorder. One interesting finding of this study was that elderly subjects had lower lifetime prevalence rates of depression and neurotic disorders than did younger groups; it is unclear whether this is a genuine cohort phenomenon,²³ or merely the result of a survivor effect or non-recall of past illness episodes by the elderly.

Not surprisingly, given the more stringent case definitions employed, the prevalence rates of neurotic disorders in epidemiological studies using the GMS/AGECAT system, a structured interview

Table 1. Prevalence and incidence rates of specific DSM-III mental disorders (ECA study)

		Males	Females	Total
<i>One-month prevalence (%)^{22,27}</i>				
Dysthymia	65+	1.0	2.3	1.8
Phobic disorder	All ages	2.2	4.2	3.3
	65+	2.9	6.1	4.8
Panic disorder	All ages	3.8	8.4	6.2
	65+	0.0	0.2	0.1
Obsessive-compulsive disorder	All ages	0.3	0.7	0.5
	65+	0.7	0.9	0.8
Somatization disorder	All ages	1.1	1.5	1.3
	65+	0.0	0.2	0.1
Generalized anxiety disorder	All ages	0.0	0.2	0.1
	65+	-	-	1.9 ^a
	45-64	-	-	3.1 ^a
<i>Annual incidence per 100 person-years of risk²¹</i>				
Phobic disorder	65+	2.66	5.52	4.29
	All ages	2.33	5.38	3.98
Panic disorder	65+	0.00	0.07	0.04
	All ages	0.30	0.76	0.56
Obsessive-compulsive disorder	65+	0.12	1.00	0.64
	All ages	0.39	0.92	0.69

^a Six-month prevalence.

women over 65 years, and after cognitive impairment higher period prevalence phobic disorder, panic disorder and somatization disorder and somatization disorder and somatization subjects with neurotic disorder before their fifties, but phobic disorder, panic disorder and somatization disorder tended to be of most neurotic disorders in both sexes, but this was phobic disorder and obsessive disorder. One interesting finding elderly subjects had lower rates of depression and neurotic disorder than younger groups; it is a genuine cohort phenomenon the result of a survivor effect of past illness episodes by the

even the more stringent case study prevalence rates of neurotic disorders using the structured interview

study)

Females	Total
2.3	1.8
4.2	3.3
6.1	4.8
8.4	6.2
0.2	0.1
0.7	0.5
0.9	0.8
1.5	1.3
0.2	0.1
0.2	0.1
-	1.9 ^a
-	3.1 ^a
5.52	4.29
5.38	3.98
0.07	0.04
0.76	0.56
1.00	0.64
0.92	0.69

with a computerized diagnostic algorithm,^{24,25} were much lower than those found by the ECA study (Table 2). However, a significant proportion had subcase levels of anxiety, phobic and obsessional neurosis. Age and sex differences for the case and subcase AGECA categories were less consistent than those found by the ECA study using DSM-III criteria. A three-year follow-up of the Liverpool sample found an incidence rate for neurotic disorders of 4.4/1000 per year.²⁶ In their view, the findings supported the idea of a general neurotic syndrome, with the majority of affected subjects having a prolonged course and variation in the predominance of different symptoms over time.

Another study of neurotic disorders in elderly community populations is the Guy's/Age Concern survey, which investigated anxiety disorders using instruments specifically developed for the study and validated against clinical diagnosis.²⁷ The survey employed nonhierarchical, nonjudgemental diagnostic criteria closely approximating in the case of phobic disorders to nonhierarchical DSM-III criteria.¹⁰ The overall one-month prevalence was 10% for phobic disorders, and 3.7% for generalized anxiety; rates were higher in women than in men, although this was only statistically significant for phobic disorders. No subject met DSM-III criteria for panic disorder. Phobic disorders in this sample were examined further in a case-control study¹⁹; cases had more neurotic symptoms than controls and higher rates of previous psychiatric disorders. One-third of the cases had their onset after the age of 65 years. Phobic

Table 2. One-month prevalence rates (%) of GMS/AGECAT neurotic disorders^{24,25}

	Male		Female	
	Case	Subcase	Case	Subcase
<i>London/New York</i>				
Anxiety neurosis	0.3	17.9	1.2	15.7
Phobic neurosis	0.0	0.6	0.0	2.1
Obsessional neurosis	0.0	5.4	0.4	4.2
Hypochondriacal neurosis	1.3	0.0	0.2	0.2
<i>Liverpool</i>				
Anxiety neurosis	0.2	18.5	1.7	16.9
Phobic neurosis	0.0	3.7	1.2	5.6
Obsessional neurosis	0.0	2.4	0.2	1.4
Hypochondriacal neurosis	0.5	0.2	0.5	0.2

subjects reported higher rates of contact with their general practitioners, but only a minority were receiving any form of treatment for their anxiety. In every case this took the form of psychotropic medication; no one had been offered any form of psychological management.

Little is known about the longer-term course and outcome of neurotic disorders in the elderly. In their review of studies of general adult populations, Marks and Lader²⁸ concluded that 41–59% of cases of 'anxiety neurosis' (generalized anxiety and panic) were recovered or much improved at 1- to 20-year follow-up. Noyes and Clancy²⁹ found that 33% of their patient sample was unchanged or worse at five-year follow-up, and that later age of onset of anxiety was a predictor of poor outcome, particularly in men. In the three-year follow-up study by Larkin *et al.*,²⁶ only 20% of the reinterviewed elderly cases had improved. In a 35-year follow-up of former panic patients Coryell³⁰ found that they had an excess mortality. In particular, male patients had a higher rate of deaths from cardiovascular disease. Follow-up studies of general adult phobic populations have shown that about half of the subjects show at least some improvement.^{31,32}

Factors associated with neurotic disorders

Biological

The biological aspects of neurotic disorders in the elderly have not yet received much attention from researchers. The evidence and its limitations have recently been reviewed by Philpot.³³ Studies of young adults show that there is a significant genetic component to vulnerability to neurotic disorders as a group,^{34,35} but it is not disorder-specific, except perhaps in obsessive-compulsive disorder (OCD) and panic disorder.^{36,37} Vulnerability factors such as fearfulness, neuroticism and intelligence are also under a degree of genetic control.^{38–41} The contribution of genetic factors to affective disorders diminishes with age, but it is not known if this is also true of neurotic disorders.

Neuroimaging studies of brain structure and function have yet to contribute significantly to our understanding of neurotic disorders in the elderly. Computerized tomography (CT) scan studies of elderly depressed patients have found that milder cases and those with higher anxiety scores are

more likely to have normal scans.^{42,43} Studies of poststroke anxiety disorders show that the distribution of lesions differs from that in patients with poststroke affective disorders, but they are not consistent as to location; for example, Sharpe *et al.*⁴⁴ found anxiety to be related to the size of left hemisphere lesions, whereas Castillo *et al.*⁴⁵ found it to be associated with right hemisphere lesions. Functional neuroimaging studies carried out with younger subjects have found changes in regional cerebral blood flow (rCBF) associated with exposure to the feared stimulus in phobic patients and with induction of obsessional thoughts in patients with OCD.⁴⁶⁻⁵⁰ Abnormalities of caudate nucleus metabolism in OCD have also been found using positron emission tomography (PET).⁵¹

While epidemiological studies suggest that there is no association between neurotic disorders and dementia, in the early stages dementia may reveal itself through symptoms such as anxiety or obsessionality.^{52,53} Anxiety in demented patients may be associated with depression or psychotic symptoms, or with the implications of the dementia and its impact on social functioning. Studies have yet to examine the possibility that the neurodegenerative process in Alzheimer's disease is itself a cause of anxiety.

Psychosocial

At all ages, there is a clear relationship between psychological ill health and indicators of social adversity such as low occupational class, unemployment, poor housing, overcrowding and limited access to amenities such as transport. In the elderly, the evidence for this association is strongest in studies that use dimensional scales for measuring anxiety and depression.^{54,55} The relationship between adversity and categorical definitions of cases is less clear; in the ECA study generalized anxiety was the only neurotic disorder associated with a socioeconomic variable (low household income). Phobic disorders in the elderly are associated with urban domicile,^{56,57} but this may not be an indicator of adversity so much as poor social networks. Adversity increases levels of distress, but is less important as an aetiological factor in the development of more severe disorders.

Adverse life events are provoking agents which determine the onset of some psychiatric disorders in vulnerable individuals.⁵⁸⁻⁶⁰ It is the individual

meaning of the event, rather than its severity, that appears to be important,⁶¹ with loss events leading to depression, and threatening events to anxiety.⁶² Some classes of life-event, such as bereavement, retirement and institutionalization, are commoner in old age than at other times of life, but their impact in relation to the onset of neurotic disorders in the elderly is still under-researched. Bereavement appears to take the same form and follow the same course in old age as it does earlier in life, with only a minority developing a psychiatric disorder.⁶³ It is not known if psychiatric morbidity following bereavement is more or less common in the elderly compared with other age groups. It has been suggested that the 'timeliness' of some losses in old age may reduce their impact^{64,65}; in one recent study, the only significant predictor of persistent grief in old age was the unexpectedness of the death.⁶⁶

Experiences such as early parental loss and physical and sexual abuse in childhood are significant vulnerability factors for depression and other psychiatric disorders in adult life, although their significance in neurotic disorders in old age has not been extensively studied. There is an association between phobic disorders, particularly agoraphobia, and early parental loss in both the young and the elderly.^{19,67,68} Generalized anxiety has also been linked with early parental loss in men.⁶⁹ It appears that it is not so much parental loss that is important, so much as associated experiences such as prior marital conflict or subsequent inadequate care.⁷⁰ Presumably these experiences affect the developing personality and result in particular cognitive habits and 'defence styles'⁷¹ that determine the responses to adverse events and experiences in later life. The relationship between abuse in childhood and psychiatric disorder in old age is not known; one small study has suggested a link between childhood sexual abuse and panic disorder in the elderly.⁷²

Physical illness

In elderly community populations, neurotic disorders are associated with increased mortality and physical morbidity.^{73,74} This association is even more marked in psychiatric and medical patients,^{75,76} which emphasizes the importance of careful history-taking and physical examination, particularly in late-onset cases.

rather than its severity, that ant,⁶¹ with loss events lead- d threatening events to anx- s of life-event, such as ent and institutionalization, age than at other times of in relation to the onset of the elderly is still under- nt appears to take the same ame course in old age as it h only a minority develop- ler.⁶³ It is not known if psy- owing bereavement is more he elderly compared with as been suggested that the sses in old age may reduce e recent study, the only sig- sistent grief in old age was the death.⁶⁶

early parental loss and use in childhood are sig- actors for depression and ers in adult life, although otic disorders in old age studied. There is an asso- c disorders, particularly parental loss in both the ^{9,67,68} Generalized anxiety ith early parental loss in t is not so much parental much as associated expe- arital conflict or subse- e.⁷⁰ Presumably these developing personality r cognitive habits and termine the responses to riences in later life. The se in childhood and psy- age is not known; one d a link between child- panic disorder in the

ulations, neurotic disor- increased mortality and his association is even iatric and medical sizes the importance of physical examination, ases.

There are several reasons why neurotic and physical disorders are linked in old age.^{77,78} Physical illness may be a cause of the neurotic disorder, either directly by presenting with neurotic symptoms (Table 3), or indirectly as an adverse life-event leading to psychological symptoms or the onset of an anxiety or depressive disorder. For example, mild, chronic anxiety symptoms are relatively common following myocardial infarction in old age,⁷⁹ and in a few cases this can develop into a severe and disabling 'cardiac neurosis', often focused on somatic anxiety symptoms such

as palpitations. Such anxiety about physical illness can have important behavioural consequences (see below).

The somatic symptoms of neurotic disorders may be mistaken for physical illness. It is a common clinical impression that the somatization of psychological distress is commoner in elderly patients,^{80,81} but to some extent this may be due to a selection bias, since it is the individuals who complain of physical symptoms of anxiety and depression who will be most likely present to medical services. Neurotic disorders may also

Table 3. *Physical causes of neurotic symptoms in the elderly*⁷⁸

<i>Cardiovascular</i>	<i>Neurological</i>
Myocardial infarction	Head injury
Cardiac arrhythmias	Cerebral tumour
Orthostatic hypotension	Dementia
Mitral valve prolapse	Delirium
	Epilepsy
	Migraine
	Cerebral lupus erythematosus
	Demyelinating disease
	Vestibular disturbance
	Subarachnoid haemorrhage
	CNS infections
<i>Respiratory</i>	<i>Dietary and drug-related</i>
Pneumonia	Caffeine
Pulmonary embolism	Vitamin deficiencies
Emphysema	Anaemia
Asthma	Sympathomimetics
Left-ventricular failure	Dopamine agonists
Hypoxia	Corticosteroids
Chronic obstructive airways disease	Withdrawal syndromes
Bronchial carcinoma	Akathisia
	Digoxin toxicity
	Activating serotonin reuptake inhibitors (e.g. fluoxetine)
<i>Endocrine and metabolic</i>	
Hypo- and hyperthyroidism	
Hypo- and hypercalcaemia	
Cushing's disease	
Carcinoid syndrome	
Hypoglycaemia	
Insulinoma	
Phaeochromocytoma	
Hyperkalaemia	
Hypokalaemia	
Hypothermia	

cause physical illness by direct or indirect effects on bodily function. Most important in this respect are the higher rates of damaging behaviours such as smoking and alcohol abuse. Physiological effects, such as anxiety-related hyperventilation causing coronary artery spasm, may also be significant.⁶²

Specific neurotic disorders

Generalized anxiety

In generalized anxiety there is persistent anxious mood accompanied by motor tension, autonomic symptoms, apprehensiveness and hypervigilance. In recent years, generalized anxiety has suffered as a diagnostic entity at the expense of the other 'specific' anxiety disorders such as phobia and panic; it has been a particular victim of the categorical approach to neurosis. Nevertheless, there remains a group of patients who are excessively and unreasonably preoccupied with general concerns without developing the symptoms and behaviours associated with other specific disorders. It has been suggested that this diagnosis may be inappropriately applied to elderly people because of their increased vulnerability and physical frailty⁸³; in fact, epidemiological evidence indicates that only a small percentage of the elderly population meet the diagnostic criteria for this disorder. In the ECA Piedmont Health Survey, the six-month prevalence of DSM-III Generalized Anxiety Disorder (GAD) without major depression or panic in elderly subjects was only 1.9%, lower than the rate found in middle-aged subjects (3.1%).⁵⁷ The prevalence rates of cases identified by other studies are similarly low.²⁴⁻²⁷ These findings suggest that old age is not associated with an inappropriate inflation of diagnoses of GAD using current classification systems; however, it is true that at the subcase level rates are much higher.^{24,25} GAD as currently defined is associated with an increased use of both physical and mental health services in the elderly.⁵⁷ If service use is regarded as a relevant criterion of clinical importance of a disorder, then the concept of generalized anxiety is still useful in this age group, particularly in primary care settings.

Phobic disorders

Phobias are defined as the persistent and irrational fear of an object, activity or situation resulting in

a compelling desire to avoid the phobic stimulus. Phobias are usually subdivided by modern classifications into agoraphobia, social phobia and simple (specific) phobia. The fears reported by elderly people are similar to those in younger age groups - animals, heights, going out of doors, public transport, etc.,¹⁹ although the relative frequency may differ; the commonest fear expressed by the urban elderly in one study was using public transport. Much is made of the 'reasonableness' of some of these fears expressed by elderly people, particularly those who live in rundown inner-city areas, and as a result clinically significant fears may be dismissed as rational. In fact, perceptions of vulnerability are determined principally by factors such as gender, physical disability and the availability of social support,⁸⁴⁻⁸⁶ and it is these rather than age that should be taken into consideration when judging the reasonableness or otherwise of any fears that are expressed.

Phobic disorders in the elderly fall broadly into two groups. On the one hand there are those with long-standing fears, usually specific in nature, which are associated with little in the way of distress or social impairment. These individuals have arranged their lives so that they need not confront their fears, and it is only occasionally, with the onset of old age, that exposure becomes unavoidable; for example, an agoraphobic may have to contend with the challenge of doing the shopping after the death of their spouse, a needle phobic with the onset of insulin-dependent diabetes, or a lift phobic with the need to use one following admission into residential care. The second important group of phobic disorders in the elderly have their onset late in life, are usually agoraphobic in nature, and are associated with clinically significant levels of distress and disability.¹⁹ These late-onset disorders typically develop following a traumatic event such as an episode of physical illness, a mugging or a fall, and the resulting impairment usually persists long after the physical consequences of the event have resolved. Panic attacks alone are rarely a cause of agoraphobia developing in old age, although they may occur as a consequence of it.

Both falls and the fear of falling are relatively common in old age, and are associated with increased levels of anxiety and depression.⁸⁷⁻⁸⁹ To some extent this association is mediated by factors such as physical disability and the prescription of psychotropic drugs. Vetter and Ford⁸⁷

to avoid the phobic stimulus. subdivided by modern classification, social phobia and simple phobia. The fears reported by elderly are similar to those in younger age groups, going out of doors, heights,¹⁹ although the relative frequency commonest fear expressed in one study was using public transport.⁸⁴ It is made of the 'reasonable' fears expressed by elderly compared with those who live in rundown areas as a result clinically dismissed as rational. In fact, disabilities are determined principally by such factors as gender, physical disability, lack of social support,⁸⁴⁻⁸⁶ rather than age that should be taken into account when judging the reasonableness of fears that are expressed. In the elderly fall broadly into two categories; on the one hand there are those with usually specific in nature, such as claustrophobia, with little or no disability. These individuals have fears that they need not confront except occasionally, with the exposure becomes unavoidable. An agoraphobic may have to change of doing the shopping without spouse, a needle phobic with insulin-dependent diabetes, or a claustrophobic who needs to use one following medical care. The second important group of disorders in the elderly have anxiety disorders are usually agoraphobic in nature associated with clinically significant physical disability.¹⁹ These later in life develop following a major illness, an episode of physical illness, and the resulting impairment long after the physical condition has resolved. Panic disorder is a cause of agoraphobia although they may occur

warn against the vicious circle of frailty, fear of falling and sedative medication, leading to further unsteadiness and falls. In some cases, fear of falling in the elderly may be associated with neurological disorder. Marks⁹⁰ has described the condition of 'space phobia', an intense fear of falling in the absence of visual supports; typically, the onset of this fear is relatively late in life, and in some individuals it is associated with disturbances of the vestibulo-ocular and righting reflexes.

Unfortunately, the psychological effects of traumatic physical health-events in old age are still poorly appreciated, with the result that both family and health and social services may unwittingly collude with the phobic avoidance by providing well intentioned but misguided help and domiciliary support. Very few elderly people with disabling phobic disorders receive any appropriate treatment for their problem²⁷; indeed, at all ages the inappropriate prescription of benzodiazepine drugs may aggravate the development of phobic symptoms.^{91,92}

Panic disorder

Panic disorder is a condition characterized by recurrent attacks of intense fear accompanied by severe somatic anxiety symptoms such as palpitations, dyspnoea and choking. Very little is known about panic in old age. Epidemiological studies indicate that panic attacks and panic disorder are rare in elderly community populations. However, it is an episodic disorder, so its prevalence is likely to be underestimated in cross-sectional surveys.⁹³ The limited evidence from case reports,^{94,95} volunteer samples⁹⁶ and nonpsychiatric patient populations^{97,98} suggests that panic in old age is less common than in early adulthood, is commoner in women and widows, and that late-onset cases are symptomatically less severe than those whose disorder started earlier in life. Elderly panic patients tend not present to psychiatric services,^{12,99} but because of the prominent physical symptoms they may be referred instead for investigation and treatment to cardiologists, neurologists and gastroenterologists. In a study of cardiology patients with chest pain and no evidence of coronary artery disease, one-third of those aged 65 years and over met diagnostic criteria for panic disorder.⁹⁸ At all ages, there is extensive comorbidity between panic disorder and somatization disorder,¹⁰⁰ and patients with panic are often misdiag-

nosed by their GPs as chronic somatizers.⁹⁷ In late-onset cases, the possibility of depression or an iatrogenic cause such as dopamine agonist treatment for Parkinson's disease should be considered.¹⁰¹ In some cases, panic attacks result in secondary agoraphobic avoidance.⁹⁴ Panic disorder has been associated with increased risk of both suicidal and nonsuicidal mortality in studies of general adult samples,^{30,102} but it is not known to what extent panic is a specific cause of increased mortality in old age.

Post-traumatic stress disorder

Exposure to extreme catastrophe can cause significant psychological disturbance, and because of the close relationship of these stress reactions to acute or ongoing trauma they are now classified as specific diagnoses in DSM-IV and ICD-10, notably post-traumatic stress disorder (PTSD). Little is known about the long-term effect of severely traumatic experiences, although PTSD can persist for many years. Its onset may also be delayed, sometimes manifesting itself for the first time in old age following retirement or an adverse life-event.^{103,104} Many people, now elderly, were exposed to severe trauma during the Second World War, and it appears that what we now call PTSD was common then and has been persistent. In one study of American ex-prisoners of war, 67% had suffered from PTSD at the time, and only 27% had fully recovered.¹⁰⁵ Another recent study of 800 World War II prisoners of war found that 80% had persistent nightmares, and that those who had been imprisoned for longer periods and who had been subjected to more severe stress were more likely to meet the diagnostic criteria for PTSD.¹⁰⁶ Traumatic wartime experiences have an enduring effect on a significant minority of older people, particularly more vulnerable groups such as psychiatric patients or those in long-term care,^{107, 108} and health professionals need to be aware of this when making assessments. Now that PTSD sufferers are eligible for financial compensation, increasing numbers of veterans are coming forward for diagnosis and treatment, and psychiatric services will need to develop better strategies for dealing with this problem.

A related issue is the nature and extent of elderly people's response to disastrous and traumatic experience in old age.¹⁰⁹ In an interesting

natural experiment, Phifer and Norris¹¹⁰ were able to examine prospectively the response of elderly inhabitants of a poor rural region of the USA to two devastating floods. They found that personal loss of property was associated with increased levels of anxiety and depression, but that this response was limited, except in individuals exposed to extreme catastrophe. Persistent psychological disturbance was related not so much to personal loss as to the degree of damage suffered by the community as a whole. Any disturbance, persistent or not, was relatively mild, and few subjects would have qualified for a DSM-III-R diagnosis of PTSD. However, some might have met the criteria for the as yet little-researched ICD-10 diagnosis of 'acute stress reaction'. The psychological resilience of many people in the face of extreme trauma and loss reminds us that we have much to learn about vulnerability and protective factors in relation to neurotic and affective disorders.

Obsessive-compulsive disorder (OCD)

People with OCD suffer from obsessive thoughts and/or compulsive acts which are a significant source of distress, or interfere with social functioning. Modern nosologies are probably most appropriate in their categorization of OCD as a distinct disorder, although its current classification with the anxiety disorders may be inappropriate. Although a proportion of OCD patients also develop significant symptoms of depression and anxiety, it is a relatively persistent and stable diagnosis.¹¹¹ OCD shows an early and specific response to treatment with serotonergic drugs,¹¹² and the placebo response rate is much lower than that which occurs in depression and anxiety.^{113,114} OCD appears to have more in common with neurodevelopmental disorders such as Gilles de la Tourette's syndrome, and Insel¹¹⁵ has suggested that it is due to impairment of a specific neuronal circuit involving the orbitofrontal cortex, basal ganglia, substantia nigra and ventrolateral pallidum. This is in contrast to other anxiety disorders, which are thought to be due to disturbances in the frontal cortex-amygdala-septum-hippocampus system which mediates fear.¹¹⁶

The clinical features of OCD in elderly patients are similar to those seen in younger adults. It is very rare for OCD to appear for the first time in old age, although cases have been reported.¹¹⁷ Most

cases that present late are long-standing disorders that have never been identified or treated.¹¹⁸ It is important therefore that all elderly cases receive thorough evaluation and the benefit of modern treatments as and when they present to psychogeriatric services.¹¹⁹ In old age, the late appearance of apparently obsessional orderliness and preoccupation with routines may presage the onset of dementia, particularly if the frontal lobe is involved. Obsessional symptoms may also appear at any age following head injury or cerebral tumour; in such cases, apparently obsessional and stereotypic behaviour is not preceded by mounting anxiety or followed by a release of tension.

Somatoform disorders

The frank somatization of psychological distress usually has its onset early in life (by definition, DSM-IV somatization disorder begins before the age of 30 years) and once established it tends to have a chronic course. Somatizing individuals are skilled both at seeking medical treatment and at avoiding psychiatrists, and it is not uncommon for these individuals to present to psychiatric services for the first time in old age. They are usually referred with a very extensive history of complaints, specialist consultations and investigations, and the clinical picture is often complicated by the presence of true physical illness, depression and anxiety. Studies of clinical populations suggest that somatization disorder does not ameliorate substantially with age,^{120,121} although as they get older these patients may transfer their concern to genuine organic symptoms. Such patients are rare, but they give rise to management problems quite out of proportion to their number.

Another form of preoccupation with physical illness more often encountered in elderly patients is hypochondriasis. In contrast to somatizing patients who have multiple complaints and who demand relief from their distress, the concern of hypochondriacal patients is typically restricted to a limited number of body systems or organs; as a rule, they are concerned with the possibility that they might be suffering from a serious physical illness, and their demand is for investigation rather than treatment. In the elderly, primary hypochondriasis is nearly always a chronic problem, and late onset hypochondriasis is usually secondary to depression or anxiety.

are long-standing disorders
identified or treated.¹¹⁸ It is
that all elderly cases receive
and the benefit of modern
when they present to psy-
.¹¹⁹ In old age, the late appear-
obsessional orderliness and
routines may presage the onset
ularly if the frontal lobe is
al symptoms may also appear
ing head injury or cerebral
es, apparently obsessional and
r is not preceded by mounting
by a release of tension.

of psychological distress
early in life (by definition,
n disorder begins before the
once established it tends to
. Somatizing individuals are
g medical treatment and at
and it is not uncommon for
resent to psychiatric services
old age. They are usually
extensive history of com-
ultations and investigations,
is often complicated by the
ical illness, depression and
clinical populations suggest
order does not ameliorate
^{120,121} although as they get
ay transfer their concern to
ms. Such patients are rare,
management problems quite
their number.

reoccupation with physical
untered in elderly patients
contrast to somatizing
multiple complaints and who
their distress, the concern of
ts is typically restricted to
dy systems or organs; as a
d with the possibility that
from a serious physical ill-
is for investigation rather
derly, primary hypochon-
a chronic problem, and
is usually secondary to

Hysterical dissociation

Hysterical symptoms in elderly patients are an important exception to the general rule that neurotic symptoms in the elderly have the same diagnostic significance as in younger adults. Apparently hysterical conversion reactions and dysmnasias occasionally appear in elderly patients in response to stress. However, as Bergmann¹²² has said, 'it is best to assert dogmatically that primary hysterical illness does not begin in old age'; such symptoms are usually due either to underlying undiagnosed physical illness, or the release of dissociative tendencies in vulnerable personalities by organic cerebral pathology or functional psychiatric disorder.

Management

Despite the availability of many behavioural, cognitive and environmental strategies for the management of neurotic disorders at all ages, drugs are still the most common form of treatment offered to elderly sufferers. Old people are the heaviest consumers of psychotropic medication, particularly benzodiazepine tranquilizers and hypnotics,¹²³ and certain groups, such as those in long-term care, are particularly likely to receive these drugs.¹²⁴ Although drug treatment is easy and convenient in the short term, it can lead to considerable problems later on, which could be avoided if the benefits and advantages of alternative approaches were more widely appreciated.

Since most cases of neurotic disorder in the elderly are seen in primary care and general medical settings, this is where the focus of identification and treatment should be. Specialist psychogeriatric services have only a limited role in the management of these patients, particularly in the long term, but they are a valuable source of expertise and should aim to offer advice and education to primary care and general hospital staff in appropriate management strategies.

Psychological treatments

Cognitive-behaviour therapy (CBT) is of proven benefit in the treatment of conditions such as phobias and OCD in younger adults, and individual case-reports and small case-series of successful treatments in elderly patients with phobic disorders¹²⁵⁻¹²⁷ show that this approach can also

be successfully applied to this age group. CBT has also been used in elderly patients with depression, prolonged grief reactions, generalized anxiety and panic with encouraging results.¹²⁸⁻¹³⁰ CBT involves both the cognitive and behavioural approaches and aims to conceptualize and modify the disturbances in thinking and behaviour that characterize neurotic disorders. Although theoretically distinct they are in practice rarely carried out in isolation, and procedures such as anxiety management training draw on both models.¹³¹ CBT may be carried out both individually and in groups. The advantages of groups are that they are more cost-effective, and they also harness useful peer support which often persists after the formal treatment has finished.¹³² However, it can be difficult and time-consuming to assemble groups in which the members are sufficiently similar to ensure they work well together and do not exclude or scapegoat particular individuals. While the principles of applying CBT to elderly patients are the same as for younger adults, the goals and techniques may need to be adapted in some cases,¹³¹ for example if there is significant physical disability. Certain behavioural strategies such as flooding which involve high levels of arousal may be less useful in the elderly, because the associated impairment in cognitive performance inhibits learning. There is no evidence that elderly patients are less likely to be able to use CBT than younger patients.

Some knowledge of psychodynamics is of great value when assessing elderly patients, particularly if their disorders involve difficulties in negotiating the developmental issues of late life, or the inappropriate and maladaptive use of defence mechanisms.¹³³ Formal psychodynamic treatments and techniques, particularly in group settings, are also useful in the treatment of neurotic disorders in elderly patients.

Drug treatments

Before starting an elderly patient on any psychotropic medication there are a number of important factors to consider. First, will it have an adverse effect on the patient's physical condition? Secondly, will the medication interact with any other drugs (including alcohol) that the patient is taking? Thirdly, what are the pharmacokinetics and pharmacodynamics of the patient? Last, but not least, is the patient likely to take the

drugs as prescribed? When making this judgement, the risk of suicide must also be taken into consideration.

Benzodiazepines There are surprisingly few formal controlled trials of benzodiazepine treatment in elderly patients, and as Salzman¹³⁴ has pointed out, those that have been published are difficult to interpret because of the very diverse nature of the subjects included and the methods used to investigate them. Current views on benzodiazepine use in this age group are based for the most part on clinical experience, extrapolation from studies of younger adults, and studies of pharmacokinetics in the elderly. Important problems associated with the prolonged use of benzodiazepines include dependence, memory impairment, poor motor coordination, depression of respiratory drive and paradoxical excitement.¹³⁵⁻¹³⁷ Elderly people are particularly sensitive to these adverse effects, and the accumulation of drugs with long elimination half-lives also leads to drowsiness, delirium, incontinence, falls and fractures. In one recent study of elderly hospitalized patients, benzodiazepines were found to account for 29% of new episodes of acute confusion.¹³⁸ As a rule, therefore, benzodiazepines in the elderly should be restricted for the most part to the management of transient, short-term anxiety symptoms, and drugs with short half-lives and no active metabolites, such as oxazepam, are to be preferred. Maintenance treatment may be appropriate in a few elderly cases, but long-term use of benzodiazepines should be avoided and established users weaned off their medication whenever possible.¹²³ Despite the problems encountered when treating elderly patients with benzodiazepines, their use is still widespread,¹³⁹ although shorter-acting compounds are now more commonly prescribed.

Antidepressants If depression is a prominent feature of the neurotic disorder, then treatment with antidepressant medication should be considered. Generalized anxiety and panic also respond to antidepressant drugs. Both tricyclics and the selective serotonin reuptake inhibitors (SSRIs) are effective, but the latter are preferable because they can be used safely in severely physically ill and suicidal patients, although activating SSRIs may actually worsen anxiety, at least in the short term (Table 3).¹⁴⁰ Drugs with serotonin-reuptake inhibiting activity also appear to have a specific

effect in OCD, and are effective in the elderly.^{119,141} Monoamine oxidase inhibitors (MAOIs) are relatively well tolerated by the elderly, although their use has declined in recent years. A specific MAO-B inhibitor, moclobemide, has recently been introduced which has much less peripheral activity than the older MAOIs, and therefore there is less need for dietary restrictions. Whether or not it has any particular role in the treatment of neurotic disorders is not known. While some neurotic symptoms may remit along with the depression following antidepressant medication, others such as established phobic avoidance will usually require a separate psychological intervention.¹⁴²

Neuroleptics The risk of disabling extrapyramidal side-effects means that neuroleptic drugs have only a limited role in the long-term management of neurotic symptoms in most elderly patients. However, short courses of these drugs are of benefit when fear and agitation are secondary to psychotic experiences, as in delirium, and in such cases haloperidol and zuclopenthixol are the drugs of choice.

Other drugs Low doses of beta-blocking drugs such as propranolol may be useful in elderly patients with prominent sympathetic somatic symptoms, although there is no published evidence of their effectiveness in this age group. They should not be used in patients with asthma, chronic obstructive airways disease, cardiac conduction defects and heart failure. These restrictions, together with side-effects such as nightmares and insomnia, limit their usefulness in the elderly.

Antihistamine drugs such as hydroxyzine have long been used as anxiolytics in the elderly. Their effect is probably due primarily to their sedative action. They are relatively safe, although hypotension can be a problem. They have a role in patients where respiratory depressant drugs are contraindicated.

Buspirone is a relatively new azapirone anxiolytic that differs both chemically and pharmacologically from the benzodiazepines. Its pharmacokinetics, safety and efficacy in the elderly are similar to those in younger adults,¹⁴³ and it appears to be well tolerated by this age group. Its short-term use is not associated with rebound, dependence or abuse.¹⁴⁴ There is no cross-tolerance with benzodiazepines, so

and are effective in the
amine oxidase inhibitors
ely well tolerated by the
use has declined in recent
D-B inhibitor, moclobemide,
duced which has much less
an the older MAOIs, and
need for dietary restrictions.
any particular role in the
disorders is not known.
symptoms may remit along
following antidepressant med-
established phobic avoid-
re a separate psychological

of disabling extrapyrami-
that neuroleptic drugs have
the long-term management
in most elderly patients.
of these drugs are of ben-
tation are secondary to psy-
in delirium, and in such
zuclopenthixol are the

es of beta-blocking drugs
may be useful in elderly
ent sympathetic somatic
ere is no published evi-
ess in this age group. They
n patients with asthma,
ays disease, cardiac con-
art failure. These restric-
effects such as nightmares
r usefulness in the elderly.
uch as hydroxyzine have
lytics in the elderly. Their
rimarily to their sedative
y safe, although hypoten-
a. They have a role in
ory depressant drugs are

ely new azapirone anxi-
hemically and pharmaco-
benzodiazepines. Its
y and efficacy in the
se in younger adults,¹⁴³
ell tolerated by this age
e is not associated with
or abuse.¹⁴⁴ There is
n benzodiazepines, so

buspirone should not be used to cover benzodi-
azepine withdrawal. Unlike other anxiolytics, it
takes about two weeks to become effective, so it
has a limited role in the management of acute
anxiety states.

References

- 1 Lindsay J. Neurotic disorders. *Rev Clin Gerontol* 1992; 2: 311-21.
- 2 World Health Organization. *International classification of diseases*, 10th revision. Geneva: World Health Organization, 1992.
- 3 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*, fourth edition. Washington: American Psychiatric Association, 1994.
- 4 Tyrer P. *Classification of neurosis*. Chichester: John Wiley, 1989.
- 5 Tyrer P. The division of neurosis: a failed classification. *J R Soc Med* 1990; 83: 614-16.
- 6 Goldberg DP, Bridges K, Duncan-Jones P *et al.* Dimensions of neurosis seen in primary care settings. *Psychol Med* 1987; 17: 461-70.
- 7 Mackinnon A, Christiansen H, Jorm AF *et al.* A latent trait analysis of an inventory designed to detect symptoms of anxiety and depression using an elderly community sample. *Psychol Med* 1994; 24: 977-86.
- 8 Goldberg D, Huxley P. *Common mental disorders: a bio-social model*. London: Tavistock/Routledge, 1992.
- 9 Lindsay J. Introduction: the concept of neurosis. In: Lindsay J ed. *Neurotic disorders in the elderly*. Oxford: Oxford University Press, 1995.
- 10 Lindsay J, Banerjee S. Phobic disorders in the elderly: a comparison of three diagnostic systems. *Int J Geriatr Psychiatry* 1993; 8: 387-93.
- 11 Office of Population Censuses and Surveys. *Mental health statistics for England*. London: HMSO, 1983.
- 12 Thyer BA, Parrish RT, Curtis GC *et al.* Ages of onset of DSM-III anxiety disorders. *Comp Psychiatry* 1985; 26: 113-22.
- 13 Schwartz GM, Braverman BG, Roth B. Anxiety disorders and psychiatric referral in the general medical emergency room. *Gen Hosp Psychiatry* 1987; 9: 87-93.
- 14 Shepherd M, Cooper B, Brown AC *et al.* *Psychiatric illness in general practice*. London: Oxford University Press, 1981.
- 15 Cooper B. Mental illness, disability and social conditions among old people in Mannheim. In: Häfner HG, Moschel N, Sartorius N eds. *Mental health and the elderly*. Berlin: Springer, 1986.
- 16 Macdonald AJD. Do general practitioners 'miss' depression in elderly patients? *BMJ* 1986; 292: 1365-67.
- 17 Morgan K, Dallosso H, Ebrahim S *et al.* Mental health and contact with primary care services in old age. *Int J Geriatr Psychiatry* 1987; 2: 223-26.
- 18 Thompson JW, Burns BJ, Bartko J *et al.* The use of ambulatory services by persons with and without phobia. *Med Care* 1988; 26: 183-98.
- 19 Lindsay J. Phobic disorder in the elderly. *Br J Psychiatry* 1991; 159: 531-41.
- 20 Robins L, Regier D eds. *Psychiatric disorders in America*. New York: Free Press, 1991.
- 21 Eaton WW, Kramer M, Anthony JC *et al.* The incidence of specific DIS/DSM-III mental disorders: data from the NIMH Epidemiologic Catchment Area program. *Acta Psychiatr Scand* 1989; 79: 163-78.
- 22 Regier DA, Boyd JH, Burke JD *et al.* One-month prevalence of mental disorders in the United States. *Arch Gen Psychiatry* 1988; 45: 977-86.
- 23 Klerman G. The current age of youthful melancholia: evidence for increase in depression among adolescents and young adults. *Br J Psychiatry* 1988; 152: 4-14.
- 24 Copeland JRM, Dewey ME, Wood N *et al.* Range of mental illness among the elderly in the community: prevalence in Liverpool using the GMS/AGECAT package. *Br J Psychiatry* 1987; 150: 815-23.
- 25 Copeland JRM, Gurland BJ, Dewey ME *et al.* The distribution of dementia, depression and neurosis in elderly men and women in an urban community: assessed using the GMS-AGECAT package. *Int J Geriatr Psychiatry* 1987; 2: 177-84.
- 26 Larkin AB, Copeland JRM, Dewey ME *et al.* The natural history of neurotic disorder in an elderly urban population. Findings from the Liverpool Longitudinal Study of Continuing Health in the Community. *Br J Psychiatry* 1992; 160: 681-86.
- 27 Lindsay J, Briggs K, Murphy E. The Guy's/Age Concern Survey: prevalence rates of cognitive impairment, depression and anxiety in an urban elderly community. *Br J Psychiatry* 1989; 155: 317-29.
- 28 Marks I, Lader M. Anxiety states (anxiety neurosis): a review. *J Nerv Ment Dis* 1973; 156: 3-18.
- 29 Noyes R, Clancy J. Anxiety neurosis: a 5-year follow-up. *J Nerv Ment Dis* 1976; 162: 200-205.

- 30 Coryell W. Mortality after thirty to forty years. Panic disorder compared with other psychiatric illnesses. In: Grinspoon L, ed. *Psychiatry update*. Washington: American Psychiatric Association, 1984.
- 31 Roberts AH. Housebound housewives - a follow-up study of a phobic anxiety state. *Br J Psychiatry* 1964; 110: 191-97.
- 32 Agras WS, Chapin N, Oliveau DC. The natural history of phobia: course and prognosis. *Arch Gen Psychiatry* 1972; 26: 315-17.
- 33 Philpot M. Biological factors. In: Lindesay J ed. *Neurotic disorders in the elderly*. Oxford: Oxford University Press, 1995.
- 34 Andrews G, Stewart G, Allen R *et al*. The genetics of six neurotic disorders: a twin study. *J Affect Disord* 1990; 19: 23-29.
- 35 Kendler KS, Kessler RC, Heath AC *et al*. Major depression and generalized anxiety disorder. Same genes, (partly) different environments? *Arch Gen Psychiatry* 1992; 49: 716-22.
- 36 Marks I. Genetics of fear and anxiety disorders. *Br J Psychiatry* 1986; 149: 406-18.
- 37 Torgersen S. Comorbidity of major depression and anxiety disorders in twin pairs. *Am J Psychiatry* 1990; 147: 1199-202.
- 38 Goldsmith HH, Gottesman II. Origins of variation in behavioural style: a longitudinal study of temperament in young twins. *Child Dev* 1981; 52: 91-103.
- 39 Bouchard TJ, Lykken DT, McGue M *et al*. The Minnesota study of twins reared apart. *Science* 1990; 250: 223-28.
- 40 Britton PG, Bergmann K, Kay DWK *et al*. Mental state, cognitive functioning, physical health and social class in the community aged. *J Gerontol* 1967; 22: 517-21.
- 41 Nunn C, Bergmann K, Britton PG *et al*. Intelligence and neurosis in old age. *Br J Psychiatry* 1974; 124: 446-52.
- 42 Jacoby R, Levy R. Computed tomography in the elderly. 3. Affective disorder. *Br J Psychiatry* 1980; 136: 270-75.
- 43 Alexopoulos GS, Young RC, Shindelacker RD. Brain computed tomography findings in geriatric depression and primary degenerative dementia. *Biol Psychiatry* 1992; 31: 591-99.
- 44 Sharpe M, Hawton K, House A *et al*. Mood disorders in long-term survivors of stroke: associations with brain lesion location and volume. *Psychol Med* 1990; 20: 815-28.
- 45 Castillo CS, Starkstein SE, Federoff JP *et al*. Generalized anxiety disorder after stroke. *J Nerv Ment Dis* 1993; 181: 102-108.
- 46 O'Carroll RE, Moffoot APR, Van Beck M *et al*. The effect of anxiety induction on the regional uptake of ^{99m}Tc -exametazime in simple phobia as shown by single photon emission tomography (SPET). *J Affect Disord* 1993; 28: 203-10.
- 47 McGuire PK, Bench CJ, Frith CD *et al*. Functional anatomy of obsessive-compulsive phenomena. *Br J Psychiatry* 1994; 164: 459-68.
- 48 Rauch SL, Jenike MA, Alpert NM *et al*. Regional cerebral blood flow measured using symptom provocation in obsessive-compulsive disorder using oxygen 15-labelled carbon dioxide and positron emission tomography. *Arch Gen Psychiatry* 1994; 51: 62-70.
- 49 Machlin SR, Harris GJ, Pearlson GD *et al*. Elevated medial-frontal cerebral blood flow in obsessive-compulsive patients: a SPECT study. *Am J Psychiatry* 1991; 148: 1240-42.
- 50 Rubin RT, Villanueva-Meyer J, Anath J *et al*. Regional xenon 133 cerebral blood flow and cerebral technetium 99m HMPAO uptake in unmedicated patients with obsessive-compulsive disorder and matched control subjects: determination by high-resolution single-photon emission computed tomography. *Arch Gen Psychiatry* 1992; 49: 695-702.
- 51 Baxter L, Schwartz J, Bergman KS *et al*. Caudate glucose metabolic rate changes with both drug and behaviour therapy for obsessive-compulsive disorder. *Arch Gen Psychiatry* 1992; 49: 681-89.
- 52 Wands K, Merskey H, Hachinski V *et al*. A questionnaire investigation of anxiety and depression in early dementia. *JAGS* 1990; 36: 535-38.
- 53 Ballard CG, Mohan RNC, Patel A *et al*. Anxiety disorder in dementia. *Ir J Psychol Med* 1994; 11: 108-109.
- 54 Himmelfarb S, Murrell SA. The prevalence and correlates of anxiety symptoms in older adults. *J Psychol* 1984; 116: 159-67.
- 55 Kennedy GJ, Kelman HR, Thomas C. Hierarchy of characteristics associated with depressive symptoms in an urban elderly sample. *Am J Psychiatry* 1989; 146: 220-22.
- 56 Walton VA, Romans-Clarkson SE, Mullen PE *et al*. The mental health of elderly women in the community. *Int J Geriatr Psychiatry* 1990; 5: 257-63.
- 57 Blazer DG, George LK, Hughes D. The epidemiology of anxiety disorders: an age comparison. In: Salzman C, Lebowitz BD eds. *Anxiety in the elderly*. New York: Springer, 1991.
- 58 Brown GW, Harris TO. *Social origins of depression*. London: Tavistock, 1978.
- 59 Brown GW, Harris TO eds. *Life events and illness*. London: Unwin Hyman, 1989.
- 60 Murphy E. Social origins of depression in old age. *Br J Psychiatry* 1982; 141: 135-42.

- single photon emission tomography. *Affect Disord* 1993; 28: 203-10.
- ench CJ, Frith CD *et al.*
- omy of obsessive-compulsive
- J Psychiatry* 1994; 164: 459-68.
- ke MA, Alpert NM *et al.*
- al blood flow measured using
- cation in obsessive-compulsive
- oxygen 15-labelled carbon
- itron emission tomography.
- Psychiatry* 1994; 51: 62-70.
- eris GJ, Pearlson GD *et al.*
- frontal cerebral blood flow in
- ulsive patients: a SPECT study.
- 1991; 148: 1240-42.
- ueva-Meyer J, Anath J *et al.*
- ¹⁵O cerebral blood flow and
- ¹⁵O HMPAO uptake in
- ents with obsessive-compulsive
- ched control subjects:
- high-resolution single-photon
- d tomography. *Arch Gen*
- 49: 695-702.
- z J, Bergman KS *et al.*
- metabolic rate changes with
- aviour therapy for obses-
- sorder. *Arch Gen Psychiatry*
- r H, Hachinski V *et al.* A
- igation of anxiety and
- dementia. *JAGS* 1990; 36:
- n RNC, Patel A *et al.*
- dementia. *Ir J Psychol Med*
- rell SA. The prevalence and
- y symptoms in older adults.
- 5: 159-67.
- n HR, Thomas C. Hierarchy
- associated with depressive
- an elderly sample. *Am J*
- 6: 220-22.
- Clarkson SE, Mullen PE *et*
- h of elderly women in the
- Geriatr Psychiatry* 1990; 5:
- K, Hughes D. The
- ety disorders: an age
- man C, Lebowitz BD eds.
- . New York: Springer,
- O. *Social origins of*
- Tavistock, 1978.
- O eds. *Life events and*
- n Hyman, 1989.
- ins of depression in old
- 1982; 141: 135-42.
- 61 Brown GW, Bifulco A, Harris TO. Life events,
- vulnerability and onset of depression: some
- refinements. *Br J Psychiatry* 1987; 150: 30-42.
- 62 Brown GW. Life events and psychiatric disorder:
- replications and limitations. *Psychosom Med*
- 1993; 55: 248-59.
- 63 Blazer DG. Late life bereavement and depressive
- neurosis. In: Blazer DG ed. *Depression in late*
- life*. New York: CV Mosby, 1982.
- 64 Zisook S, Shuchter SR, Lyons LE. Predictors of
- psychological reactions during the early stages
- of widowhood. *Psychiatr Clin N Am* 1987; 10:
- 355-68.
- 65 Davis A. Life events in the normal elderly. In:
- Copeland JRM, Abou-Saleh MT, Blazer DG eds.
- Principles and practice of geriatric psychiatry*.
- Chichester: Wiley, 1994.
- 66 Byrne G, Raphael B. A longitudinal study of
- bereavement phenomena in recently widowed
- elderly men. *Psychol Med* 1994; 24: 411-21.
- 67 Faravelli C, Webb T, Ambonetti A *et al.*
- Prevalence of traumatic early life events in 31
- agoraphobic patients with panic attacks. *Am J*
- Psychiatry* 1985; 142: 1493-94.
- 68 Tweed JL, Schoenbach VJ, George LK *et al.* The
- effects of childhood parental death and divorce
- on six-month history of anxiety disorders. *Br J*
- Psychiatry* 1989; 154: 823-28.
- 69 Zahner GER, Murphy JM. Loss in childhood:
- anxiety in adulthood. *Comp Psychiatry* 1989;
- 30: 553-63.
- 70 Tennant C, Bebbington P, Hurry J. Social
- experiences in childhood and adult psychiatric
- morbidity: a multiple regression analysis.
- Psychol Med* 1982; 12: 321-27.
- 71 Pollock C, Andrews G. The defense style
- associated with specific anxiety disorders. *Am J*
- Psychiatry* 1989; 146: 455-60.
- 72 Sheikh JI, Swales PJ, Kravitz J *et al.* Childhood
- abuse history in older women with panic
- disorder. *Am J Geriatr Psychiatry* 1994; 2:
- 75-77.
- 73 Kay DWK, Bergmann K. Physical disability and
- mental health in old age. *J Psychosom Res*
- 1966; 10: 3-12.
- 74 Lindesay J. The Guy's/Age Concern Survey:
- physical health and psychiatric disorder in an
- urban elderly community. *Int J Geriatr*
- Psychiatry* 1990; 5: 171-78.
- 75 Bergmann K. The neuroses of old age. In: Kay
- DWK, Walk A eds. *Recent developments in*
- psychogeriatrics*. Ashford: Headley Bros, 1971.
- 76 Burn WK, Davies KN, McKenzie FR *et al.* The
- prevalence of psychiatric illness in acute
- geriatric admissions. *Int J Geriatr Psychiatry*
- 1993; 8: 175-80.
- 77 Goldberg DP. Discussion of Prof. Ottoman's
- paper. In: Ohmann R, Freeman HL, Frank
- Holmkvist A, Neilzen S eds. *Interaction*
- between mental and physical illness*. Berlin:
- Springer-Verlag, 1989.
- 78 Pitt B. Neurotic disorders and physical illness.
- In: Lindesay J ed. *Neurotic disorders in the*
- elderly*. Oxford: Oxford University Press, 1995.
- 79 Peach H, Pathy J. Disability of the elderly after
- myocardial infarction. *J R Coll Physicians* 1979;
- 13: 154-57.
- 80 De Alarcon RD. Hypochondriasis and depres-
- sion in the aged. *Gerontology* 1964; 6: 266-77.
- 81 Blazer D, George L, Landerman R. The
- phenomenology of late life depression. In:
- Bebbington P, Jacoby R eds. *Psychiatric*
- disorders in the elderly*. London: Mental Health
- Foundation, 1986.
- 82 Haines AP, Imeson JD, Meade TW. Phobic
- anxiety and ischaemic heart disease. *BMJ* 1987;
- 295: 297-99.
- 83 Shamoian CA. What is anxiety in the elderly?
- In: Salzman C, Lebowitz BD eds. *Anxiety in the*
- elderly*. New York: Springer, 1991.
- 84 Hindelang MJ, Gottfredson MR, Garofalo J.
- Victims of personal crime: an empirical*
- foundation for a theory of personal victimiza-*
- tion*. Cambridge, MA: Balinger, 1978.
- 85 Yin P. Fear of crime as a problem for the
- elderly. *Soc Prob* 1982; 30: 240-45.
- 86 Fattah EA, Sacco VF. *Crime and victimization*
- in the elderly*. New York: Springer-Verlag, 1989.
- 87 Vetter NJ, Ford D. Anxiety and depression in
- elderly fallers. *Int J Geriatr Psychiatry* 1989; 4:
- 159-64.
- 88 Downton JH, Andrews K. Postural disturbance
- and psychological symptoms amongst elderly
- people living at home. *Int J Geriatr Psychiatry*
- 1990; 5: 93-98.
- 89 Arfken CL, Lach HW, Birge SJ *et al.* The
- prevalence and correlates of fear of falling in
- elderly persons living in the community. *Am J*
- Publ Health* 1994; 84: 565-70.
- 90 Marks I. Space phobia: syndrome or agorapho-
- bic variant. *J Neurol Neurosurg Psychiatry*
- 1981; 44: 387-91.
- 91 Ashton H. Benzodiazepine withdrawal: an
- unfinished story. *BMJ* 1984; 288: 1135-40.
- 92 Cohen SI. Phobic disorders and benzodiazepines
- in the elderly. *Br J Psychiatry* 1992; 160: 135.
- 93 Von Korff MR, Eaton WW. Epidemiological
- findings on panic. In: Baker R ed. *Panic*
- disorder: theory, research and therapy*. London:
- Wiley, 1989.
- 94 Luchins DJ, Rose RP. Late-life onset of panic
- disorder with agoraphobia in three patients. *Am*
- J Psychiatry* 1989; 146: 920-21.
- 95 Frances A, Flaherty JA. Elderly widow develops

- panic attacks, followed by depression. *Hosp Community Psychiatry* 1989; 40: 19-23.
- 96 Sheikh JI, King RJ, Barr Taylor C. Comparative phenomenology of early-onset versus late-onset panic attacks: a pilot survey. *Am J Psychiatry* 1991; 148: 1231-33.
 - 97 Katon WJ. Chest pain, cardiac disease and panic disorder. *J Clin Psychiatry* 1984; 51: 27-30.
 - 98 Beitman BD, Kushner M, Grossberg GT. Late onset panic disorder: evidence from a study of patients with chest pain and normal cardiac evaluations. *Int J Psychiatr Med* 1991; 21: 29-35.
 - 99 Kenardy J, Oei TPS, Evans L. Neuroticism and age of onset for agoraphobia with panic attacks. *J Behav Ther Exp Psychiatry* 1990; 21: 193-97.
 - 100 Boyd JH, Burke JD, Gruenberg E *et al.* Exclusion criteria of DSM-III: a study of hierarchy-free syndromes. *Arch Gen Psychiatry* 1984; 41: 983-99.
 - 101 Schiffer RB, Kurlan R, Rubin A *et al.* Evidence for atypical depression in Parkinson's disease. *Am J Psychiatry* 1988; 145: 1020-22.
 - 102 Weissmann MM, Klerman GL, Markowitz JS *et al.* Suicidal ideation and suicide attempts in panic disorders and attacks. *N Engl J Med* 1989; 321: 1209-14.
 - 103 Scaturo DJ, Hyman PM. The impact of combat trauma across the family life cycle: clinical considerations. *J Trauma Stress* 1992; 5: 273-88.
 - 104 Kaup BA, Ruskin PE, Nyman G. Significant life events and PTSD in elderly World War II veterans. *Am J Psychiatry* 1994; 151: 239-43.
 - 105 Kluznik JC, Speed N, Van Valkenberger C *et al.* Forty-year follow-up of United States prisoners of war. *Am J Psychiatry* 1986; 143: 1443-45.
 - 106 Guerrero J, Crocq M. Sleep disorder in the elderly: depression and PTSD. *J Psychosom Res* 1994; 38 (suppl 1): 141-50.
 - 107 Rosen J, Fields RB, Hand AM *et al.* Concurrent posttraumatic stress disorder in psychogeriatric patients. *J Geriatr Psychiatr Neurol* 1989; 2: 65-69.
 - 108 Hermann N, Eryavec G. Posttraumatic stress disorder in institutionalized World War II veterans. *Am J Psychiatry* 1994; 151: 324-31.
 - 109 McFarlane A. Life events and psychiatric disorder: the role of natural disaster. *Br J Psychiatry* 1987; 151: 362-67.
 - 110 Phifer JE, Norris FH. Psychological symptoms in older adults following natural disaster: nature, timing, duration and course. *J Gerontol* 1989; 44: S207-S217.
 - 111 Rasmussen SA, Tsuang MT. Clinical characteristics and family history in DSM-III obsessive compulsive disorder. *Am J Psychiatry* 1986; 143: 317-22.
 - 112 Goodman WK, Price LH, Rasmussen SA *et al.* Efficacy of fluvoxamine in obsessive-compulsive disorder. A double-blind comparison with placebo. *Arch Gen Psychiatry* 1989; 46: 36-44.
 - 113 Montgomery SA. Clomipramine in obsessional neurosis: a placebo-controlled trial. *Pharm Med* 1980; 1: 189-92.
 - 114 de Veauh Geiss J, Landau P, Katz R. Treatment of obsessive compulsive disorder with clomipramine. *Psychiatr Ann* 1989; 19: 97-101.
 - 115 Insel TR. Neurobiology of obsessive-compulsive disorder: a review. *Int Clin Psychopharmacol* 1992; 7 (suppl 1): 31-33.
 - 116 Gray J. *The neuropsychology of anxiety*. Oxford: Oxford University Press, 1982.
 - 117 Bajulaye R, Addonizio C. Obsessive-compulsive disorder arising in a 75-year-old woman. *Int J Geriatr Psychiatry* 1992; 7: 139-42.
 - 118 Jenike MA. *Geriatric psychiatry and psychopharmacology: a clinical approach*. St Louis, MO: CV Mosby, 1989.
 - 119 Austin LS, Zealberg JJ, Lydiard RB. Three cases of pharmacotherapy of obsessive-compulsive disorder in the elderly. *J Nerv Ment Dis* 1992; 179: 634-35.
 - 120 Guze SB, Woodruff RA, Clayton PJ. Sex, age and the diagnosis of hysteria (Briquet's syndrome). *Am J Psychiatry* 1972; 129: 745-48.
 - 121 Pribor EF, Smith DS, Yutzy SH. Somatization disorder in elderly patients. *Am J Geriatr Psychiatry* 1994; 2: 109-17.
 - 122 Bergmann K. Neurosis and personality disorder in old age. In: Isaacs AD, Post F eds. *Studies in geriatric psychiatry*. Chichester: Wiley, 1978.
 - 123 Higgitt A. Indications for benzodiazepine prescriptions in the elderly. *Int J Geriatr Psychiatry* 1992; 3: 239-49.
 - 124 Gilbert A, Quintrell LN, Owen N. Use of benzodiazepines among residents of aged-care accommodation. *Community Health Stud* 1988; 12: 394-99.
 - 125 Thyer BA. Prolonged *in-vivo* exposure therapy with a 70-year old woman. *J Behav Ther Exp Psychiatry* 1981; 12: 69-71.
 - 126 Woods RT, Britton PG. *Clinical psychology with the elderly*. London: Croom Helm/Chapman & Hall, 1985.
 - 127 Downs AFD, Rosenthal TL, Lichstein KL. Modelling therapies reduce avoidance of bath-time by the institutionalized elderly. *Behav Ther* 1988; 19: 359-68.
 - 128 Thompson LW, Gallagher DE, Breckenridge JS. Comparative effectiveness of psychotherapies for depressed elderly. *J Consult Clin Psychol* 1987;

- story in DSM-III obsessive
er. *Am J Psychiatry* 1986;
- ice LH, Rasmussen SA *et al.*
amine in obsessive-compulsive
-blind comparison with
Psychiatry 1989; 46: 36-44.
Clomipramine in obsessional
-controlled trial. *Pharm Med*
- Landau P, Katz R.
sive compulsive disorder with
Psychiatr Ann 1989; 19: 97-101.
logy of obsessive-compulsive
Int Clin Psychopharmacol
31-33.
psychology of anxiety.
iversity Press, 1982.
izio C. Obsessive-compulsive
75-year-old woman. *Int J*
1992; 7: 139-42.
c psychiatry and psy-
clinical approach. St Louis,
89.
JJ, Lydiard RB. Three cases
of obsessive-compulsive
ly. *J Nerv Ment Dis* 1992;
- RA, Clayton PJ. Sex, age
hysteria (Briquer's
Psychiatry 1972; 129: 745-48.
Yutzy SH. Somatization
tients. *Am J Geriatr*
09-17.
s and personality disorder
AD, Post F eds. *Studies in*
hichester: Wiley, 1978.
for benzodiazepine
derly. *Int J Geriatr*
39-49.
N, Owen N. Use of
g residents of aged-care
Community Health Stud 1988;
- n-vivo exposure therapy
man. *J Behav Ther Exp*
9-71.
Clinical psychology with
room Helm/Chapman &
TL, Lichstein KL.
uce avoidance of bath-
lized elderly. *Behav Ther*
- er DE, Breckenridge JS.
ss of psychotherapies for
sult Clin Psychol 1987;
- 55: 385-90.
- 129 Kavanagh DJ. Towards a cognitive-behavioural intervention for adult grief reactions. *Br J Psychiatry* 1990; 157: 373-83.
 - 130 King P, Barrowclough C. A clinical pilot study of cognitive-behavioural therapy for anxiety disorders in the elderly. *Behav Psychother* 1991; 19: 337-45.
 - 131 Woods RT. Psychological treatments I: behavioural and cognitive approaches. In: Lindsay J ed. *Neurotic disorders in the elderly*. Oxford: Oxford University Press, 1995.
 - 132 Zerhusen JD, Boyle K, Wilson W. Out of the darkness: group cognitive therapy for the elderly. *J Psychosoc Nursing* 1991; 29: 16-20.
 - 133 Martindale B. Psychological treatments II: psychodynamic approaches. In: Lindsay J. ed. *Neurotic disorders in the elderly*. Oxford: Oxford University Press, 1995.
 - 134 Salzman C. Pharmacologic treatment of the anxious elderly patient. In: Salzman C, Lebowitz BD eds. *Anxiety in the elderly*. New York: Springer, 1991.
 - 135 Tyrer P. Dependence on benzodiazepines. *Br J Psychiatry* 1980; 137: 576-77.
 - 136 Curran HV. Tranquillizing memories: a review of the effects of benzodiazepines on human memory. *Biol Psychiatry* 1986; 23: 179-213.
 - 137 Fancourt G, Castleden M. The use of benzodiazepines with particular reference to the elderly. *Br J Hosp Med* 1986; 5: 321-25.
 - 138 Foy A, O'Connell D, Henry D *et al.* Benzodiazepine use as a cause of cognitive impairment in elderly hospital inpatients. *J Gerontol* 1995; 50: M99-M106.
 - 139 Stewart R, Marits RG, Padgett PD *et al.* Benzodiazepine use in an ambulatory elderly population. *Clin Ther* 1994; 16: 118-24.
 - 140 Evans ME, Lye M. Depression in the physically ill; an open study of treatment with the 5-HT reuptake inhibitor fluoxetine. *J Clin Exp Gerontol* 1992; 14: 297-307.
 - 141 Jenike MA. *Handbook of geriatric psychopharmacology*. Littleton, MA: PSG Publishing, 1985.
 - 142 Blazer DG, Hughes DC, Fowler N. Anxiety as an outcome symptom of depression in elderly and middle-aged adults. *Int J Geriatr Psychiatry* 1989; 4: 273-78.
 - 143 Robinson D, Napoliello MJ, Shenck L. The safety and usefulness of buspirone as an anxiolytic drug in elderly versus young patients. *Clin Ther* 1988; 10: 740-46.
 - 144 Lader M. Can buspirone induce rebound, dependence or abuse? *Br J Psychiatry* 1991; 159 (suppl 12): 45-51.

